VOL. 562, NOS. 1+2 JOURNAL OF CHROMATOGRAPHY JANUARY 2, 1991

*(Biomedical Applications, Vol. 100, Nos. 1+2)*

# CONTENTS

## SECOND INTERNATIONAL SYMPOSIUM ON APPLIED MASS SPECTROMETRY IN THE HEALTH SCIENCES. BARCELONA, APRIL 17–20, 1990

Foreword
by E. Gelpi (Barcelona, Spain), A. Mallet (London, U.K.) and W. J. Richter (Basle, Switzerland) . . . . . 1

*New developments and fundamental studies*

Thermospray liquid chromatography–mass spectrometry with a quadrupole ion trap mass spectrometer
by R. E. Kaiser, Jr., J. D. Williams, S. A. Lammert and R. G. Cooks (West Lafayette, IN, U.S.A.) and D. Zakett (Midland, MI, U.S.A.) . . . . . 3

Particle beam liquid chromatography–mass spectrometry on a double-focusing sector instrument
by L. Baczynskyj (Kalamazoo, MI, U.S.A.) . . . . . 13

Thermospray and particle beam liquid chromatographic–mass spectrometric analysis of coumarin anticoagulants
by J. X. De Vries (Heidelberg, F.R.G.) and K. A. Kymber (Manchester, U.K.) . . . . . 31

Use of particle beam liquid chromatography–electron impact mass spectrometry for structure elucidation of oxodipine and three of its metabolites
by C. Julien-Larose, P. Voirin, C. Mas-Chamberlin and A. Dufour (Chilly Mazarin, France) 39

Biomedical applications of high-performance liquid chromatography–mass spectrometry with continuous-flow fast atom bombardment
by D. S. Millington, D. L. Norwood and N. Kodo (Durham, NC, U.S.A.), R. Moore (Belfast, U.K.) and M. D. Green and J. Berman (Research Triangle Park, NC, U.S.A.) . . . 47

Analysis of some eicosanoids by continuous-flow fast atom bombardment mass spectrometry
by U. Justesen and G. Bojesen (Odense, Denmark) . . . . . 59

Microbore liquid chromatography coupled to a fast atom bombardment probe for the on-line detection of the Tyr-Pro cleavage of a nonapeptide by recombinant HIV-1 protease
by S. M. Cole, P. V. Macrae, J. R. Merson, F. S. Pullen and D. J. Rance (Sandwich, U.K.) 67

Chromatographic developments in low flow delivery for liquid chromatography–mass spectrometry
by M. P. Balogh and C. C. Stacey (Milford, MA, U.S.A.) . . . . . 73

Chloride attachment negative-ion mass spectra of sugars by combined liquid chromatography and atmospheric pressure chemical ionization mass spectrometry
by Y. Kato and Y. Numajiri (Ibaraki-ken, Japan) . . . . . 81

Scanning thin-layer chromatography–liquid secondary ion mass spectrometry and its application for investigation of drug metabolites
by Y. Nakagawa and K. Iwatani (Osaka, Japan) . . . . . 99

*Clinical and metabolic studies*

Analysis of rat brain microdialysate by gas chromatography–high-resolution selected-ion monitoring mass spectrometry
by S. Y. Chang, T. A. Moore, L. L. Devaud, L. C. E. Taylor and E. B. Hollingsworth (Research Triangle Park, NC, U.S.A.) . . . . . 111

Fragmentation pathways of O-trimethylsilyl ethers of hydroxy long-chain bases analysed by linked-scan mass spectrometry
by T. Matsubara and A. Hayashi (Higashi-osaka, Japan) . . . . . 119

Automated screening of urine samples for carbohydrates, organic and amino acids after treatment with urease
by J. D. Shoemaker and W. H. Elliott (St. Louis, MO, U.S.A.) . . . . . 125

Abnormal fatty acid metabolism in patients in hopantenate therapy during clinical episodes
by M. Matsumoto, T. Kuhara, Y. Inoue, T. Shinka and I. Matsumoto (Ishikawa, Japan) . 139

Determination of very-long-chain fatty acids in plasma by a simplified gas chromatographic–mass spectrometric procedure
by U. Caruso (Genova, Italy), B. Fowler and M. Erceg (Manchester, U.K.) and C. Romano (Genova, Italy) . . . . . 147

Characteristic fragmentation of thromboxane $B_3$ in thermospray high-performance liquid chromatography–mass spectrometry
by J. Abian and E. Gelpi (Barcelona, Spain) . . . . . 153

Measurement of leucine and α-ketoisocaproic acid fluxes in the fetal/placental unit
by G. L. Loy, A. N. Quick, Jr., F. C. Battaglia, G. Meschia and P. V. Fennessey (Denver, CO, U.S.A.) . . . . . 169

Electrospray mass spectrometry in the clinical diagnosis of variant hemoglobins
by C. H. L. Shackleton (Oakland, CA, U.S.A.), A. M. Falick (San Francisco, CA, U.S.A.), B. N. Green (Altrincham, U.K.) and H. E. Witkowska (Oakland, CA, U.S.A.) . . . . . 175

Application of liquid chromatography–thermospray mass spectrometry in the analysis of glycerophospholipid molecular species
by F. A. Kuypers, P. Bütikofer and C. H. L. Shackleton (Oakland, CA, U.S.A.) . . . . 191

Mass spectral characterization of urinary metabolites of D,L-kawain
by C. Köppel and J. Tenczer (Berlin, F.R.G.) . . . . . 207

Phenylalanine in whole blood and plasma — a candidate reference method
by G. W. Lynes and M. Hjelm (London, U.K.) . . . . . 213

Detection of ribose-methylated nucleotides in enzymatic hydrolysates of RNA by thermospray liquid chromatography–mass spectrometry
by N. Takeda, S. C. Pomerantz and J. A. McCloskey (Salt Lake City, UT, U.S.A.) . . . 225

Gas chromatographic–mass spectrometric characterisation of some novel hydroxyeisosatetraenoic acids formed on incubation of arachidonic acid with microsomes from induced rat livers
by R. A. Clare (Beckenham, U.K.), S. Huang (Guildford, U.K.), M. V. Doig (Beckenham, U.K.) and G. G. Gibson (Guildford, U.K.) . . . . . 237

Use of high-performance liquid chromatography–thermospray mass spectrometry and gas chromatography–electron-impact mass spectrometry in the identification of the metabolites of α-methylacetohydroxamic acids, potential anti-asthmatic agents
by P. M. Woollard, J. A. Salmon and A. D. Padfield (Beckenham, U.K.) . . . . . 249

Characterization of phosphatidylcholines in rabbit lung lavage fluid by positive and negative ion fast-atom bombardment mass spectrometry
by H. Gasser, W. Strohmaier, G. Schlag, E. R. Schmid and G. Allmaier (Vienna, Austria) 257

Study of human urinary metabolism of fenfluramine using gas chromatography–mass spectrometry
by R. Brownsill, D. Wallace, A. Taylor and B. Campbell (Fulmer, U.K.) . . . . . 267

Isatin (indole-2,3-dione) in urine and tissues. Detection and determination by gas chromatography–mass spectrometry
by J. M. Halket, P. J. Watkins, A. Przyborowska, B. L. Goodwin, A. Clow, V. Glover and M. Sandler (London, U.K.) . . . . . 279

Mass spectrometric structure determination of metabolites of 3-[2-(N,N-dimethylaminomethyl)-phenylthio]phenol
by M. Ryska, I. Koruna, L. Poláková and E. Svátek (Prague, Czechoslovakia) . . . . . 289

Urinary metabolites of cannabidiol in dog, rat and man and their identification by gas chromatography–mass spectrometry
by D. J. Harvey (Oxford, U.K.) and E. Samara and R. Mechoulam (Jerusalem, Israel) . . 299

Studies on anabolic steroids. V. Sequential reduction of methandienone and structurally related steroid A-ring substituents in humans: gas chromatographic–mass spectrometric study of the corresponding urinary metabolites
by R. Massé, H. Bi, C. Ayotte, P. Du, H. Gélinas and R. Dugal (Pointe-Claire, Canada) . 323

On-line measurements of $^{13}C$ enrichments in rat breath. Non-invasive method for *in vivo* study of drug enzymatic induction
by R. Guilluy, F. Billion-Rey and J. L. Brazier (Lyon, France) . . . . . 341

Mass spectrometric and high-performance liquid chromatographic studies of medroxyprogesterone acetate metabolites in human plasma
by G. Sturm, H. Häberlein, T. Bauer, T. Plaum and D. J. Stalker (Marburg, F.R.G.) . . 351

Reversible reaction via a carbanion intermediate in the elimination of ammonia from L-histidine catalysed by histidine ammonia-lyase
by T. Furuta, H. Takahashi and Y. Kasuya (Tokyo, Japan) . . . . . 363

Fast atom bombardment mass spectrometry as a rapid means of screening mixtures of ether-linked polar lipids from extremely halophilic archaebacteria for the presence of novel chemical structures
by K.-D. Klöppel and H. L. Fredrickson (Braunsweig, F.R.G.) . . . . . 369

*Biopolymers*

Structural analysis of biologically active peptides and recombinant proteins and their modified counterparts by mass spectrometry
by B. N. Pramanik, A. Tsarbopoulos, J. E. Labdon, P. P. Trotta and T. L. Nagabhushan (Bloomfield, NJ, U.S.A.) . . . . . 377

Primary sequences of insulin-like growth factors 1 and 2 isolated from porcine plasma
by S. Bayne (Gentofte, Denmark), C. M. Hoogerbrugge (Utrecht, The Netherlands), J. Thomsen and L. Skriver (Gentofte, Denmark) and S. C. van Buul-Offers and J. L. van den Brande (Utrecht, The Netherlands) . . . . . 391

Mass spectrometric determination of glycosylation sites and oligosaccharide composition of insect-expressed mouse interleukin-3
by M. Svoboda and M. Przybylski (Konstanz, F.R.G.), J. Schreurs and A. Miyajima (Palo Alto, CA, U.S.A.) and K. Hogeland and M. Deinzer (Corvallis, OR, U.S.A.) . . . . . 403

Liquid secondary ion mass spectrometry applied to structural confirmation of enzymically prepared C-terminal-truncated derivatives of recombinant hirudin
by M. Maftouh, C. Ponthus, A. Tuong and C. Picard (Toulouse, France) . . . . . 421

Mass spectral characterization of C-glycosidic flavonoids isolated from a medicinal plant (*Passiflora incarnata*)
by L. Qimin, H. van den Heuvel, O. Delorenzo, J. Corthout, L. A. C. Pieters, A. J. Vlietinck and M. Claes (Wilrijk-Antwerp, Belgium) . . . . . 435

Determination of the enantiomeric purity of synthetic peptides by gas chromatography–mass spectrometry
by C. Celma and E. Giralt (Barcelona, Spain) . . . . . 447

Analysis of tyrosine- and methionine-containing neuropeptides by fast atom bombardment mass spectrometry
by J. Silberring and F. Nyberg (Uppsala, Sweden) . . . . . 459

*Environmental and toxicological studies*

Recent methods for the determination of volatile and non-volatile organic compounds in natural and drinking water
by A. Bruchet and M. F. Legrand (Le Pecq, France) and P. Arpino and D. Dilettato (Paris, France) . . . . . 469

Gas chromatographic–mass spectrometric characterization of volatile organic compounds in Barcelona tap water
by A. Guardiola, F. Ventura, L. Matia, J. Caixach and J. Rivera (Barcelona, Spain) . . . 481

Gas chromatographic–mass spectrometric analysis of urban-related aquatic and airborne volatile organic compounds. Study of the extracts obtained by water closed-loop stripping and air adsorption with charcoal and polyurethane foam
by A. Rosell, J. I. Gomez-Belinchon and J. O. Grimalt (Barcelona, Spain) . . . . . 493

Mobile phase variations in thermospray liquid chromatography–mass spectrometry of pesticides
by G. Durand, N. de Bertrand and D. Barceló (Barcelona, Spain) . . . . . 507

Analysis of triallate residues in cereals and soil by gas chromatography with ion-trap detection
by C. Sanchez-Brunete, T. Salto, J. M. Garcia-Baudin and J. L. Tadeo (Madrid, Spain) . 525

Gas chromatographic–mass spectrometric analysis of tar compounds formed during pyrolysis of rice husks
by V. A. C. Haanappel, T. W. Stevens, A. Hovestad, V. Skolnik and R. Visser (Enschede, The Netherlands) . . . . . 531

Confirmation method for the identification and determination of some organophosphorus and organochlorine pesticides in cocoa beans by gas chromatography–mass spectrometry
by P. Rodríguez and J. Permanyer (Barcelona, Spain) and J. M. Grases and C. González (Cabrils, Spain) . . . . . 547

Identification of trichothecenes by thermospray, plasmaspray and dynamic fast-atom bombardment liquid chromatography–mass spectrometry
by R. Kostiainen (Helsinki, Finland) . . . . . 555

Gas chromatographic–mass spectrometric analysis of diols and tetrols from reactions of polycyclic aromatic hydrocarbon epoxides with hemoglobin
by B. W. Day, S. Naylor, L.-S. Gan, Y. Sahali, T. T. Nguyen, P. L. Skipper, J. S. Wishnok and S. R. Tannenbaum (Cambridge, MA, U.S.A.) . . . . . 563

*Quantitative applications*

Analysis of methionine enkephalin in human pituitary by multi-dimensional reversed-phase high-performance liquid chromatography, radioreceptor assay, radioimmunoassay, fast atom bombardment mass spectrometry, and mass spectrometry–mass spectrometry
by J. L. Lovelace, J. J. Kusmierz and D. M. Desiderio (Memphis, TN, U.S.A.) . . . . . 573

Biogenic amines: their occurrence, biosynthesis and metabolism in the locust, *Schistocerca gregaria*, by gas chromatography–negative-ion chemical ionisation mass spectrometry
by R. G. Macfarlane, J. M. Midgley and D. G. Watson (Glasgow, U.K.) and P. D. Evans (Cambridge, U.K.) . . . . . 585

Determination of 6-keto-$PGF_{1\alpha}$, 2,3-dinor-6-keto-$PGF_{1\alpha}$, thromboxane $B_2$, 2,3-dinor-thromboxane $B_2$, $PGE_2$, $PGD_2$ and $PGF_{2\alpha}$ in human urine by gas chromatography–negative ion chemical ionization mass spectrometry
by C. Weber and M. Höller (Neuss, F.R.G.), J. Beetens and F. De Clerck (Beerse, Belgium) and F. Tegtmeier (Neuss, F.R.G.) . . . . . 599

Novel derivatization and immunoextraction to improve microanalysis of 11-dehydrothromboxane $B_2$ in human urine
by M. Ishibashi and K. Watanabe (Tokyo, Japan), Y. Ohyama and M. Mizugaki (Sendai, Japan), Y. Hayashi (Tokushima, Japan) and W. Takasaki (Tokyo, Japan) . . . . . 613

Microderivatization of 4-[$^{18}$O]hydroxyproline and quantitation with a benchtop mass spectrometer
by C. D. Warner and S. K. Branch (Cambridge, MA, U.S.A.), J. F. Burke and L. M. Chambers (Boston, MA, U.S.A.) and V. R. Young (Cambridge, MA, U.S.A.) . . . . . 625

Determination of N-acetylcysteine in human plasma by gas chromatography–mass spectrometry
by A. Longo, M. Di Toro, C. Galimberti and A. Carenzi (Milan, Italy) . . . . . 639

*Gas chromatography and liquid chromatography combined with mass spectrometry*

Applications of gas chromatography–mass spectrometry in the study of androgen and odorous 16-adrostene metabolism by human axillary bacteria
by A. I. Mallett (London, U.K.), K. T. Holland and P. J. Rennie (Leeds, U.K.) and W. J. Watkins and D. B. Gower (London, U.K.) . . . . . 647

Gas chromatography–mass spectrometry and gas chromatography–tandem mass spectrometry of cyclic fatty acid monomers isolated from heated fats
by J. L. Le Quéré and J. L. Sébédio (Dijon, France), R. Henry (Villeurbanne, France) and F. Couderc, N. Demont and J. C. Promé (Toulouse, France) . . . . . 659

Derivatization and gas chromatographic–mass spectrometric detection of anabolic steroid residues isolated from edible muscle tissues
by E. Daeseleire, A. De Guesquière and C. Van Peteghem (Ghent, Belgium) . . . . . 673

Gas chromatographic–mass spectrometric analysis of some potential toxicants amongst volatile compounds emitted during large-scale thermal degradation of poly(acrylonitrile–butadiene–styrene) plastic
by M. M. Shapi and A. Hesso (Helsinki, Finland) . . . . . 681

Pre-column derivatization of free bile acids for high-performance liquid chromatographic and gas chromatographic–mass spectrometric analysis
by H. E. C. S. Ferreira (Lisbon, Portugal) and W. H. Elliott (St. Louis, MO, U.S.A.) . . 697

High-performance liquid chromatographic–mass spectrometric assay of high-value compounds for pharmaceutical use from plant cell tissue culture: *Cinchona* alkaloids
by S. Abdulrahman, M. E. Harrison, K. J. Welham, M. A. Baldwin, J. D. Phillipson and M. F. Roberts (London, U.K.) . . . . . 713

Approach to the analysis of diuretics and masking agents by high-performance liquid chromatography–mass spectrometry in doping control
by R. Ventura (Barcelona, Spain), D. Fraisse, M. Becchi and O. Paisse (Vernaison, France) and J. Segura (Barcelona, Spain) . . . . . 723

Analysis of tropane alkaloids with thermospray high-performance liquid chromatography–mass spectrometry
by S. Auriola and A. Matinsen (Kuopio, Finland), K.-M. Oksman-Caldentey (Helsinki, Finland) and T. Naaranlahti (Kuopio, Finland) . . . . . 737

Abstracts of the non-chromatographic papers published in *Analytica Chimica Acta*, Vol. 241, No. 2 (1990) . . . . . 745

*Author Index* . . . . . 757

*Compound Index* . . . . . 763

In articles with more than one author, the name of the author to whom correspondence should be addressed is indicated in the article heading by a 6-pointed asterisk (*)